

AD\_\_\_\_\_

Award Number: **W81XWH-08-1-0446**

TITLE: **Role of BRCA2 in the expressions of IRF9-regulated genes in human breast cells**

PRINCIPAL INVESTIGATOR: **Gautam Chaudhuri, PhD**

CONTRACTING ORGANIZATION: **Meharry Medical College  
Nashville, TN 37208**

REPORT DATE: July 2010

TYPE OF REPORT: **Annual**

PREPARED FOR: **U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012**

DISTRIBUTION STATEMENT:

**x Approved for public release; distribution unlimited**


The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE 07-31-2010		2. REPORT TYPE Annual		3. DATES COVERED 01 July 2009 30 June 2010	
4. TITLE AND SUBTITLE <b>Role of BRCA2 in the expressions of IRF9-regulated genes in human breast cells</b>				5a. CONTRACT NUMBER W81XWH-08-1-0446	
				5b. GRANT NUMBER BC074591 IDEA Award	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) <b>Gautam Chaudhuri</b>  gchaudhuri@mmc.edu				5d. PROJECT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Meharry Medical College Nashville, TN 37208				5f. WORK UNIT NUMBER	
				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We hypothesize that BRCA2 facilitates the formation and/or the function of the ternary ISGF3 complex and thus, functional BRCA2 protein is essential for the antiproliferative effects of type I interferons against human breast tumor cells. <b>Specific aims:</b> (A) To evaluate further the structural and functional interactions of BRCA2 with the members of the ISGF3 complex (STAT1, STAT2 and IRF9) in the human breast cells. (B) To evaluate the antiproliferative effects of BRCA2 over expression in the human breast cells with or without knock down of the IRF9 protein by RNA interference. (C) To evaluate the antiproliferative effects of type I interferons against tumors developed by BRCA2 positive and BRCA2 negative human breast tumor cells in the nude mice xenograft model.					
15. SUBJECT TERMS Breast cancer, BRCA2, Interferon alpha, Interferon beta, STAT1, STAT2, IRF9, ISGF3					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  UU	18. NUMBER OF PAGES  13	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

## Table of Contents

	<u>Page</u>
Introduction.....	4
Body.....	5
Key Research Accomplishments.....	11
Reportable Outcomes.....	11
Conclusion.....	12
References.....	12
Appendices.....	13

BRCA2, a tumor suppressor whose inactivation is associated with hereditary breast and ovarian cancer predisposition, is essential for DNA repair in mammalian cells [1-5]. BRCA2-deficient cells are defective in the repair of DNA double-strand breaks by error-free homologous recombination [5, 6], allowing error-prone repair processes to create gross chromosomal re-arrangements that may promote carcinogenesis [6]. The role of BRCA2 in homologous recombination has been linked to its functions in the regulation of RAD51, a RecA-related recombinase that forms the nucleoprotein filaments on damaged DNA that are crucial to recombinational repair [7, 8]. BRCA2 binds directly to RAD51 through 6 of the 8 BRC repeats, ~30 amino acid motifs encoded



We found that transient ablation of the breast cancer susceptibility gene BRCA2 in the human breast cells (Fig. 1) impairs the expressions of many type I interferon regulated genes (Table 1). Thus, it appears that type I interferons need functional BRCA2 for their actions [14]. Thus, negative regulation of IFN-induced genes in BRCA2-ablated breast cells may reflect another growth regulatory role of BRCA2.

4

subunits, IFNAR-1 and -2, which are associated with TYK-2 and JAK-1, respectively [17-19]. TYK-2 and JAK-1 phosphorylate tyrosine residues on the receptor that provide docking sites for the src-homology-2 (SH2) domains of STATs in a cell type specific manner [15, 16]. Once phosphorylated, STATs are released from the receptor and form heterodimers. In response to Type I IFNs, STAT2 is recruited to the IFNAR1 chain, where it is phosphorylated by TYK-2 and serves as a lure for STAT1 [15, 16]. Once released from the receptor, the resulting STAT1:STAT2 heterodimer associates with IRF9, a DNA binding protein (also called p48), forming a complex named IFN-stimulated gene factor-3 (ISGF3). After formation, ISGF3 translocates to the nucleus where it binds to the IFN-stimulated response elements (ISRE) upstream of IFN response genes and initiates transcription (Fig. 1).

**We hypothesize** that BRCA2 facilitates the formation and/or the function of the ternary ISGF3 complex and thus, functional BRCA2 protein is essential for the antiproliferative effects of type I interferons against human breast tumor cells.

**Specific aims to verify the hypothesis are:** **(A)** To evaluate further the structural and functional interactions of BRCA2 with the members of the ISGF3 complex (STAT1, STAT2 and IRF9) in the human breast cells. **(B)** To evaluate the antiproliferative effects of BRCA2 over expression in the human breast cells with or without knock down of the IRF9 protein by RNA interference. **(C)** To evaluate the antiproliferative effects of type I interferons against tumors developed by BRCA2 positive and BRCA2 negative human breast tumor cells in the nude mice xenograft model.

## **BODY**

**Task outlined in the approved Statement of Work for this period of the project**

### **Task#1**

**To evaluate further the structural and functional interactions of BRCA2 with the members of the ISGF3 complex (STAT1, STAT2 and IRF9) in the human breast cells. (Months 1-18)**

- (a) Evaluation of requirements of the BRCA2 domains in the BRCA2/ISGF3 binding by pull down assays using three recombinant fragments (N-terminal, middle, and C-terminal of approximately equal lengths) of human BRCA2 protein linked to GST-tag. (Months 11-15).
- (b) Further evaluation of direct binding between BRCA2 and ISGF3 components by yeast 2-hybrid analysis. (Months 6-18)

### **Task#2**

**To evaluate the antiproliferative effects of BRCA2 over expression in the human breast cells with or without knock down of the IRF9 protein by RNA interference. (Months 16-30)**

- (a) Generation of MDA-MB-231 and BT-549 cell derivatives inducibly over expressing BRCA2 and their characterization. (Months 16-19).

- (b) Knock down of IRF9 gene expression in the control and the BRCA2-over expressing cells. (Months 19-22).
- (c) Evaluation of the levels of BRCA2 and IRF9 mRNAs and proteins in the cells by RT-PCR and Western blotting, respectively. (Months 22-23)

**1. We evaluated the ability of different domains of human BRCA2 protein in pulling down the components of the BRCA2/ISGF3 complex.** Human BRCA2 is a large protein (3418 amino acids). We have amplified 5 consecutive fragments from the human BRCA2 ORF (in plasmid pCINBRCA2WT). There are five peptide fragments: the first 4 fragments are from the N-terminal end and have 683 amino acids (ORF ~2055 bp with start and stop codons) and the 5<sup>th</sup> C-terminal fragment has 686 amino acids. We have cloned these PCR amplified products into N-terminal GST-tagged protein expression vector [pFN2K (GST) Flexi® Vector]. We expressed these N-terminal GST-tagged BRCA2 peptide fragments in human breast cells (Fig. 2). By GST-pull down assays we found that only the N-terminal 683 amino acid fragment of BRCA2 was able to pull down all three components of the ISGF3 complex (Fig. 2). We conclude that BRCA2 interacts with the ISGF3 complex perhaps through the N-terminal domains. As it does not interact with its BRC repeats with ISGF3, ISGF3 will not compete with RAD51.

#### BRCA2-F1

MPIGSKERPTFFEIFKTRCNKADLGPISLWFEELSSEAPPYNSEPAEESSEHKNNNYEPNLFKTPQRKPSYNQLASTPI  
IFKEQGLTLPLYQSPVKELDKFKLDLGRNVPNSRHKSLRTVKTMDQADDVSCPLLNSCLSESPVVLQCTHVTPQRDK  
SVVCGSLFHTPKFVKGRQTPKHISESLGAEVDPDMSWSSSLATPPTLSSTVLIVRNEEASETVFPHDTTANVKSYSFN  
HDESLKKNDRFIASVTDSSENTNQREAASHGFGKTSNGSFVNSCKDHIGKSMPNVLEDEVYETVVDTSEEDSFSLCF  
SKCRTKNLQKVRTSKTRKKIFHEANADECEKSKNQVKEKYSFVSEVEPNDDPLDSNVANQKPFESGSDKISKEVVP  
SLACEWSQLTSLGLNGAQMEKIPLLHSSCDQNISEKDLLDTENKRKKDFLTSENLSLPRISSLPKSEKPLNEETVVNKR  
DEEQHLESHTDCILAVKQAIISGTSPVASSFQGIKKSIFRIRESKETFNASFSGHMTDPNFKKETEASESGLEIHTVCSQ  
KEDSLCPNLIDNGSWPATTTQNSVALKNAGLISTLKKKTNKFIYAIHDETSYKGKKIPKDKSELINCSAQFEANAFEAP  
LTFANADSGLLHSSVKRSCSQNDSEEP TSLTSSFGTILRKCSRNCTCSNNTVIS

#### BRCA2-F2

QDL DYKEAKCNKEKLQLFITPEADSLSCLQEGQCENDPKSKKVSDIKEEVLAACHPVQHSKVEYS DTD FQS QKSLLY  
DHENASTLILPTSKDVLSNLVMISRGKESYKMSDKLKGNNYESDVELTKNIPMEKNQDVCALNENYKNVELLPPEKY  
MRVASPSRKVQFNQNTNLRVIQKNQEETTSISKITVNPDSSEELFSDNENNFVFQVANERNNLALGNTKELHETDLTCV  
NEPIFKNSTMVLYGDTGDKQATQVSIKKDLVYVLAENKNSVKQHIKMTLGQDLKSDISLNDIKIPEKNNDYMNKWAGL  
LGPISNHSFGGSFRITASNKEIKLSEHNKSKMFFKDIEEQYPTSLACVEIVNTLALDNQKKLSKPQSINTVSAHLQSSV  
VVDCKNSHITPQMLFSKQDFNSNHNLTSPQKAEITELSTILEESGSQFEFTQFRKPSYILQKSTFEVPENQMTILKTTT  
EECDADLHVIMNAPSIGQVDSSKQFEGTVEIKRKFAGLLKND CNKSASGYLTDENEVGFGRGFYSAHGTKLNVSTEAL  
QKAVKLFSDIENISEETSAEVHPISLSSSKCHDSVSMFKIENHNDKTVSEKNNKCQLILQNNIEMTTGTFVEEITENYK  
RNTENEDNKYTAASRNSHNLEFDGSDSSKNDTVCIHKDETDLFTDQHNICL

#### BRCA2-F3

KLSGQFMKEGNTQIKEDLSDLTFLEVAKAQEACHGNTSNKEQLTATKTEQNIKDFETSDTFFQTASGKNISVAKESFN  
KIVNFFDQKPEELHNFSLNSELHSDIRKNKMDILSYEETDIVKHKILKESVPVGTGNQLVTFQGPQPERDEKIKEPTLLGF  
HTASGKKVKIAKESLDKVKNLDFDEKEQGTSEITSFSHQWAKTLKYREACKDLELACETIEITAAPKCKEMQNSLNNDKN  
LVSIETVVPKLLSDNLCRQTENLKTSSKIFLKVKVHENVETAKSPATCYTNQSPYSVIENSALAFYTSCSRKTSVVSQ  
TSLEAKKWLRGIFDQGPERINTADYVGNLYENNSNSTIAENDKNHLSEKQD TYLSNSSMSNSYSYHSDEYVND  
GYLSKNKLD SGIEPVLKNVEDQKNTSFSKVISNVKDANAYPQTVNEDICVEELVTSSSPCKNKNAIKLSISNSNNFEV  
GPPAFRIASGKIVCVSHETIKKVKDIFTDSFSKVIKENNENKSKICQTKIMAGCYEALDDSEDILHNSLDNDECSTHSHKV  
FADIQSEEILQHNQNM SGLKVS KISPCDVSLETSDICKCSIGKLHKS VSSANTCGIFSTASGKSVQVSDASLQNA RQV  
FSEIEDSTKQVFSKVLFKSNEHSDQLTREENTAIRTPHEHLISQKGFSYNV


#### BRCA2-F4

VNSSAFSGFSTASGKQVSILESSLHKVKGVL EEFDLIRTEHSLHYSPTS RQNVSKILPRVDKRNPEHCNVNSEMEKTCS  
KEFKLSNNLNVEGGSENHNSIKVSPYLSQFQQDKQQLVLGTVSLVENIHVLGKEQASPKNVKMEIGKTETFS DVPV  
KTNIEVCSTYSKDSENYFETEAVEIAKAFMEDDELTD SKLP SHATHSLFTCPENEEMVLSNSRIGKRRGEPLILVGEPSI

KRNLLNEFDRIIENQEKSLKASKSTPDGTIKDRRLFMHHVSLEPITCVPFRTTKERQEIQNPNFTAPGQEFLSKSHLYEH  
 LTLEKSSSNLAVSGHPFYQVSATRNEKMRHLITGRPTKVFPFVKTSKSHFHRVEQCVRNINLEENRQKQNIIDGHGSD  
 DSKNKINDNEIHQFNKNNSNQAAAVTFTKCEEEPLDLITSLQNARDIQDMRIKKKQRQVRFPQPGSLYLAKTSTLPRISL  
 KAAVGGQVPSACSHKQLYTYGVSKHCIKINSKNAESFQFHTEDYFGKESLWTGKGIQLADGGWLIPSNDBGKAGKEEF  
 YRALCDTPGVDPKLISRIWVYNHYRWIWKLAAMECAFPKEFANRCLSPERVLLQLKYRYDTEIDRSRRSAIKKIMERD  
 DTAAKTLVLCVSDIISLANISETSSNKTSSADTQKVAIIEITDGWYAVKAQL

### BRCA2-F5

DPPLLAVLKNRGLTVGQKIILHGAELVGSPDACTPLEAPESLMLKISANSTRPARWYTKLGFFPDPRPFPLPLSSLFSD  
 GGNVGCVDVIIQRAYPIQWMEKTSSGLYIFRNEREEEEKEAAKYVEAQQKRLEALFTKIQEEFEEHEENTTKPYLPSRAL  
 TRQQVRALQDGAELYEAVKNAADPAYLEGYFSEEQLRALNNHRQMLNDKKQAQIQLEIRKAMESAEQKEQGLSRDV  
 TTVWKLRIVSYSKKEKDSVLSIWRPSSDLYSLLTEGKRYRIYHLATSKSKSKSERANIQLAATKKTQYQQLPVSDIILF  
 QIYQPREPLHFSKFLDPDFQPSCSEVDLIGFVVSVKKTGLAPFVYLSDECYNLLAIKFWIDLNEDIKPHMLIAASNQW  
 RPESKSGLLTLFAGDFSVFSASPKEGHFQETFNKMKNTVENIDILCNEAENKLMHILHANDPKWSTPTKDCSTGPYTA  
 QIIPGTGNKLLMSSPNCEIYYQSPLSLCMAKRKSVSTPVSAQMTSKSCKGEKEIDDQKNCKKRRALDFLSRLPLPPP  
 SPICTFVSPAAQKAFQPPRSCGTYETPIKKKELNSPQMTPFKKFNEISLLESNSIADEELALINTQALLSGSTGEKQFIS  
 VSESTRTAPTSSDYLRLLKRRCTTSLIKEQESSQASTEECEKNKQDTITTKKYI




**Fig. 1. Amino acid sequences of different fragments of human BRCA2 protein tested for their abilities to bind to the components of the ISGF3 complex.**

**2. Further evaluation of direct binding between BRCA2 and ISGF3 components by yeast 2-hybrid analysis.** We employed BD Clontech Matchmaker Gold yeast two hybrid system for this purpose. We cloned the five BRCA2 ORF fragments described in the previous section into pGBKT7 DNA-BD cloning vector. Similarly we have cloned IRF9 and Stat-2 (wild type and Y690A mutant) ORFs into pGADT7 AD cloning vector. We are also cloned Stat-1 ORFs into pGADT7 vector. The yeast cells were co-transfected individually with pGADT7 fusion construct and pGBKT7 fusion construct. Yeast cells were plated onto synthetic dropout medium lacking leucine, tryptophan, and histidine in the presence of 5-bromo-4-chloro-3-indolyl-ft-D-galactopyranoside (X-α-Gal; Clontech) to select for yeast containing weaker interacting proteins. Yeast cells were also plated onto synthetic dropout medium lacking leucine, tryptophan, histidine, and adenine in the presence of 5-bromo-4-chloro-3-indolyl-ft-D-galactopyranoside (X-α-Gal; Clontech) to select for yeast containing stronger interacting proteins. The positive control (supplied with the reagent kit) used was SV40 T-antigen and p53, known to



interact very strongly. The negative controls used were cells co-transfected with empty pGADT7/pGBKT7 vector, untransformed AH109 cells and singly transformed yeast cells.  $\beta$ -Galactosidase assays were done using the Beta Glo Assay reagents and protocols (Promega) to detect  $\beta$ -Galactosidase activity in the cotransfected yeast cell extracts. Yeast cells as well as the Beta-Glo reagent were brought to room temperature.




Equal volumes of the reagent and the yeast cell culture were added together. The sample contents were mixed for 30 seconds. Samples were then incubated for 30 min at room temperature and the luciferase activity was measured using a luminometer. Both the cell growth data (not shown) as well as the beta-glo assay data (Fig. 3) suggest that none of the five BRCA2 fragments directly binds to the individual component proteins of the ISGF3 complex in the yeast cells. These data may suggest that BRCA2 perhaps interacts with the ISGF3 complex once it is formed as a trimer.

**Fig. 3. Beta-glo assay showing no direct binding of any of the five BRCA2 protein fragments with IRF9, STAT1 or STAT2 individually inside the yeast cells.**

**3. Generation of MDA-MB-231 and BT-549 cell derivatives inducibly over expressing BRCA2 and their characterization.** We have generated lentiviral constructs for full length BRCA2 with a C-terminal FLAG tag, as described before for SLUG [20, 21]. We transfected MDA-MB-231 and BT549 cells constitutively expressing tet-repressor protein with this construct, selected for stable transfectants and then evaluated the mRNA and protein levels of recombinant BRCA2 in the presence or absence of doxycycline. We evaluated recombinant BRCA2 mRNA levels with primers designed from BRCA2 ORF and the FLAG epitope. These primers did not amplify the native BRCA2 in the cells. We performed real-time RT-PCR analysis for this evaluation. Total RNA was isolated from the cultured cells using TRIzol reagent (Invitrogen). The cDNA was synthesized from 1  $\mu$ g of total RNA using the iScript cDNA Synthesis kit (BioRad). Real-time PCR quantification was performed following standard protocols using Syber green dye (BioRad). RT-PCR was performed in the iCycler (BioRad): 95 °C for 10 min, 40 cycles of 15 s at 95 °C, 30 s at 51 °C, 30 s at 72 °C followed by 1 min at 95 °C, 30 s at 55 °C and 30 s at 95 °C. The fold change over control samples was calculated using  $CT$ ,  $\Delta CT$ , and  $\Delta\Delta CT$  values [20, 21].  $\beta$ -Actin RNA was used as an endogenous control. For the evaluation of BRCA2 protein levels in the breast cancer cell lines by Western blotting with FLAG antibody, protein bands were developed using IR Dye 800 conjugated secondary antibody (LI-COR Biosciences), and visualized using LI-COR's Odyssey Infrared Imaging System. Quantitation and analysis of bands were performed using Odyssey's software.  $\beta$ -actin was used as normalization control. Figs. 4A and 4B






**Fig. 4. Over expression of FLAG-tagged BRCA2 in MDA-MB-231 and BT549 cells.** (A) Evaluation of recombinant BRCA2 mRNA levels by real-time RT-PCR. (B) Evaluation of recombinant BRCA2 protein by quantitative immunoblot analysis using FLAG antibody. Cells not treated with doxycycline were used as corresponding control. Results are mean  $\pm$  SEM (n=6). The differences were statistically significant  $p < 0.001$ .

show the over expression of recombinant BRCA2 mRNA and protein in these stably transfected cells.


**4. Knockdown of IRF9 gene expression in the control and the BRCA2-over expressing cells.** IRF9 siRNAs and corresponding control siRNAs were designed using the Block-IT RNAi designer software (Invitrogen) and purchased from Invitrogen. The nucleotide sequences of these siRNAs and respective control RNAs used in this study are as follows: Stealth961: 5'-GAGCUCUUCAGAACCGCCUACUUCU-3'/5'-AGAAGUAGGCGGUUCUGAAGAGCUC-3'; Control961: 5'-GAGUCCUGAAACCCGUCCAUUCUCU-3'/5'-AGAGAAUGGACGGGUUUCAGGACUC-3'; Stealth1025: 5'-CACCGAAGUCCAGGUAACACUGAA-3'/5'-UUCAGUGUUACCUGGAACUUCGGUG-3'; Control1025: 5'-CACGAACUUGACAUGACAUCCGGAA-3'/5'-UCCGGAUGUCAUGUCAAGUUCGUG-3'.

Transfection of these siRNAs into the breast cells was done by lipofection using the Lipofectamine 2000 (Invitrogen) as per the manufacturer's instructions. Briefly, cells were transfected at ~50% confluence using 100 pmol siRNA in six well plates and whole-cell lysates were prepared 48 h after transfection. We isolated RNA from these cells using Trizol reagents (Invitrogen). Knockdown of the expressions of the target mRNAs by the experimental siRNA and the corresponding protein were verified by real-time RT-PCR and immunoblot analysis, respectively [20, 21] as described in the previous section. Our data (Fig. 5A and 5B) shows the successful knockdown of IRF9 mRNA (Fig. 5A) and protein (Fig. 5B) in the MDA-MB-231 and BT549 cells with the Stealth siRNAs used.



**Fig. 5. Knockdown of IRF9 in MDA-MB-231 and BT549 cells using two different Stealth siRNAs.** (A) Evaluation of IRF9 mRNA levels by real-time RT-PCR. (B) Evaluation of IRF9 protein by quantitative immunoblot analysis. Cells treated with control siRNA were used as corresponding control. Results are mean  $\pm$  SEM (n=6). The differences were statistically significant  $p < 0.001$ .

Fig. 6 shows the inhibition of the ISG15 gene promoter activity in the IRF9 knocked down cells. We evaluated ISG15 gene promoter activity in transiently transfected cells by dual luciferase assay [20, 21].



We PCR amplified human ISG15 gene promoter from total DNA isolated from MDA-MB-231 cells with specific primers. The amplified DNA was cloned into the pCR4.0/TOPO vector (Invitrogen) and subsequently subcloned into the Hind III/Pst I sites of pRL-Null vector (Promega). Cells were seeded on 24-well tissue culture plates in triplicate and allowed to grow overnight to reach 90% to 95% confluence. The following day cells were transfected with pGL3-Control and pRL-ISG15 promoter construct using Lipofectamine

2000 transfection reagent (Invitrogen). Forty-eight hours later, luciferase activities were measured using the Dual Luciferase Reporter Assay System (Promega) [20, 21]. *Renilla* Luciferase activity was normalized to firefly luciferase activity [20, 21].

### Ongoing experiments:

- (i) Determination of cell growth rate by thymidine incorporation assay, by colorimetric assay, clonogenic assay and flow cytometric analysis.
- (ii) Evaluation of the tumorigenicity of the BRCA2 over expressing cells with or without knock down of IRF9.

### Key Research Accomplishments

- N-terminal sequences of human BRCA2 protein appears to be involved in the direct binding of this protein with the ISGF3 protein complex as was revealed by pull down analysis.
- None of the fragments of BRCA2 protein could directly bind with individual components of the ISGF3 complex in the yeast 2-hybrid analysis, suggesting that BRCA2 perhaps binds with the ternary complex after it is formed.
- We were successful in over expressing FLAG-tagged BRCA2 protein in an inducible manner in MDA-MB-231 and BT549 cells.
- We successfully knocked down IRF9 in the breast cancer cells.
- Knock down of IRF9 negatively affected ISG15 gene promoter activity in the knocked down cells.

**REPORTABLE OUTCOMES:** We have not yet published or presented the research performed in this grant project. But the research performed in this project directly or indirectly contributed to the following publications and poster abstracts.

### Publication:

1. Mittal, M. K., Myers, J. N., Bailey, C. K., Misra, S. and **Chaudhuri, G.** (2010) Mode of action of the retrogene product SNAILP, a SNAIL homolog, in human breast cancer cells. ***Mol. Biol Report* 37**, 1221-1227.
2. Misra, S. Sharma, S., Agarwal, A., Khedkar, S. V., Tripathi, M. K., Mittal M. K., and **Chaudhuri, G.** (2010) Cell cycle-dependent regulation of the bi-directional overlapping promoter of human BRCA2/ZAR2 genes in breast cancer cells. ***Molecular Cancer* 9**, 50.

### Meeting abstracts:

The current research on this project directly or indirectly affected the studies performed in the following poster presentations from our lab during 2009-2010:

1. Mittal, M. K., Singh, K. and **Chaudhuri, G.** (2009) Mechanisms of SLUG-induced drug resistance development in breast cancer cells. **Presented as poster at the 2009 San Antonio Breast Cancer Symposium (SABCS) held in Henry B. Gonzalez Convention Center, San Antonio, Texas, USA, December 9-13, 2009.**
2. Bailey, C. K., Mittal, Mukul, Misra, Smita and **Chaudhuri, G.** (2010) Reduction of the invasive phenotype of SNAIL-over expressing human breast cancer cells by peptide aptamer-mediated inhibition of SNAIL protein functions. **Presented at the AACR Annual meeting in Washington DC on April 17-21, 2010.**

3. Mittal, M. K., and **Chaudhuri, G.** (2010) Repression of alpha-, beta- and gamma-catenin gene expressions by SNAI2 in human breast cancer cells. **Presented at the AACR Annual meeting in Washington DC on April 17-21, 2010.**
4. Hall, Mack III, Misra, Smita, and **Chaudhuri, G.** (2010) Molecular analysis of the physical interactions of *Trypanosoma brucei* BRCA2 with different RAD51 isoforms. **Presented at the ASBMB Annual meeting in Anaheim, CA on April 24-28, 2010.**
5. Misra, Smita, and **Chaudhuri, G.** (2010) Regulation of BRCA2 gene expression through CpG methylation of its bi-directional promoter induced by endogenous siRNAs. **Presented at the ASBMB Annual meeting in Anaheim, CA on April 24-28, 2010.**

### **Conclusion:**

BRCA2 interacts with the ISGF3 ternary complex through its N-terminal domain. We are currently evaluating the effect of IRF9 knockdown in breast cells on the tumor suppressive effects of BRCA2 protein.

### **REFERENCES:**

- [1] Pestka, S., Langer, J. A., Zoon, K. C., and Samuel, C. E. (1987) Interferons and their actions. *Annu. Rev. Biochem.* **56**, 727-777.
- [2] Vilcek, J. (2006). Fifty years of interferon research: aiming at a moving target. *Immunity* **25**, 343–348.
- [3] Pestka, S., Krause, C. D., and Walter, M. R. (2004). Interferons, interferonlike cytokines, and their receptors. *Immunol. Rev.* **202**, 8–32.
- [4] van Boxel-Dezaire, A. H. H., Rani, M. R. S., and Stark, G. R. (2006). Complex modulation of cell type-specific signaling in response to type I interferons. *Immunity* **25**, 361–372.
- [5] Takaoka, A., and Yanai, H. (2006) Interferon signalling network in innate defense. *Cell. Microbiol.* **8**, 907-922.
- [6] Imai, K., and Takaoka, A. (2006) Comparing antibody and small-molecule therapies for cancer. *Nat. Rev. Cancer.* **6**, 714-727.
- [7] Nicolini, A., Carpi, A., and Rossi, G. (2006) Cytokines in breast cancer. *Cytokine Growth Factor Rev.* **17**, 325-337.
- [8] Toma, S., Raffo, P., Nicolo, G., Canavese, G., Margallo, E., Vecchio, C., Dastoli, G., Iacona, I., and Regazzi-Bonora, M. (2000) Biological activity of all-transretinoic acid with and without tamoxifen and alpha-interferon 2a in breast cancer patients. *Int. J. Oncol.* **17**, 991-1000.
- [9] Recchia, F., Frati, L., Rea, S., Torchio, P., and Sica, G. (1998) Minimal residual disease in metastatic breast cancer: treatment with IFN-beta, retinoids, and tamoxifen. *J. Interferon Cytokine Res.* **18**, 41-47.
- [10] Kornek, G., Reiner, A., Sagaster, P., Stierer, M., Mayer, A., and Ludwig, H. (1999) Effect of interferon alpha-2a on hormone receptor status in patients with advanced breast cancer. *Cancer Invest.* **17**, 189–194.

- [11] Tripathi, M. K., and Chaudhuri, G. (2005) Down-regulation of UCRP and UBE2L6 in BRCA2 knocked-down human breast cells. *Biochem. Biophys. Res. Commun.* **328**, 43-48.
- [12] de Veer, M. J., Holko, M., Frevel, M., Walker, E., Der, S., Paranjape, J. M., Silverman, R. H., and Williams, B. R. (2001) Functional classification of interferon-stimulated genes identified using microarrays. *J. Leukoc. Biol.* **69**, 912-920.
- [13] Shamoo, Y. (2003) Structural insights into BRCA2 function. *Curr. Opin. Struct. Biol.* **13**, 206-211.
- [14] Shivji, M. K., and Venkitaraman, A. R. (2004) DNA recombination, chromosomal stability and carcinogenesis: insights into the role of BRCA2. *DNA Repair (Amst)*. **3**, 835-843.
- [15] Turner, N., Tutt, A., and Ashworth, A. (2004) Hallmarks of 'BRCAness' in sporadic cancers. *Nat. Rev. Cancer.* **4**, 814-819.
- [16] Rudkin, T. M., and Foulkes, W. D. (2005) BRCA2: breaks, mistakes and failed separations. *Trends Mol. Med.* **11**, 145-148.
- [17] Pellegrini, L., and Venkitaraman, A. (2004) Emerging functions of BRCA2 in DNA recombination. *Trends Biochem. Sci.* **29**, 310-316.
- [18] Rahman, N., and Stratton, M. R. (1998) The genetics of breast cancer susceptibility. *Annu. Rev. Genet.* **32**, 95-121.
- [19] Powell, S. N., Willers, H., and Xia, F. (2002) BRCA2 Keeps Rad51 in line. high-fidelity homologous recombination prevents breast and ovarian cancer? *Mol. Cell* **10**, 1262-1263.
- [20] Tripathi, M. K., Misra, S., Khedkar, S. V., Hamilton, N., Irvin-Wilson, C., Sharan, C., Sealy, L., and Chaudhuri, G. (2005) Regulation of BRCA2 gene expression by the SLUG repressor protein in human breast cells. *J. Biol. Chem.* **280**, 17163-17171.
- [21] Mittal, M., Myers, J. N., Misra, S., Bailey, C. K. and Chaudhuri, G. (2008) *In vivo* binding to and functional repression of the VDR gene promoter by SLUG in human breast cells. *Biochem. Biophys. Res. Commun.* **372**, 30-34.

APPENDICES: None.